

FIG. 1A:ICV/P recommendations 2020-21.

Vaccine	Age in completed weeks/months/years																
	Birth	6w	10w	14w	6m	7m	9m	12m	13m	15m	16-18m	18-24m	2-3 Y	4-6 Y	9-14 Y	15-18 Y	
BCG																	
Hepatitis B	HB 1 ^a	HB 2	HB 3	HB 4 ^b													
Polio	IPV 1 ^c	IPV 2 ^c	IPV 3 ^c								IPV BL			IPV B2			
DTPa/DTP	DPT 1	DPT 2	DPT 3								DPT 01			DPT 02			
Hib	Hib 1	Hib 2	Hib 3	Hib 3							Hib BL						
PCV	PCV 1	PCV 2	PCV 3							PCV E							
Rotavirus	RV 1	RV 2	RV 3 ^d														
Influenza					Dose 1 ^e	Dose 2					Annual Vaccination						
MMR					Dose 1		Dose 2				Dose 2			Dose 3			
TCV																	
Hepatitis A								Dose 1						Dose 2 ^f			
Varicella														Dose 1			
Tdap ^g /Td																	
HPV																	
Meningococcal ^h																	
JE								Dose 1	Dose 2								
Cholera								Dose 1	Dose 2								
PPSV 23								Dose 1	Dose 2								
Rabies																	
Yellow Fever																	

Recommended age

Catch up age range

Vaccines in special situations

- To be given within 24 h after birth. When this is missed, it can be administered at first contact with health facility.
- An extra dose of Hepatitis B vaccine is permitted as part of a combination vaccine when use of this combination vaccine is necessary.
- IPV can be given as part of a combination vaccine.
- 3rd dose of Rota vaccine is not necessary for RV1.
- Influenza vaccine should be started after 6 mo, 2 doses 4 wks apart, usually in the pre-monsoon period. At other times of the year, the most recent available strain should be used. Annual influenza vaccination should be continued, for all, till 5 y of age.
- Single dose is to be given for the live attenuated Hepatitis A vaccine. The inactivated vaccine needs two doses.
- 2nd dose of Varicella vaccine should be given 3-6 mo after dose 1. However, it can be administered anytime 3 mo after dose 1 or at 4-6 y.
- Tdap should not be administered as the second booster of DTP at 4-6 y. For delayed 2nd booster, Tdap can be given after 7 y of age. A dose of Tdap is necessary at 10-12 y, irrespective of previous Tdap administration. If Tdap is unavailable/unaffordable, it can be substituted with Td.
- Before 14 completed years, HPV vaccines are recommended as a 2-dose schedule, 6 mo apart.
- From 15th y onwards and the immunocompromised subjects at all ages, HPV vaccines are recommended as a 3-dose schedule, 0-1-6 (HPV2) or 0-2-6 (HPV4)
- Menactrais approved in a 2-dose schedule between 9-23 mo. Minimum interval between two doses should be 3 mo. Menveois recommended as a single dose schedule after 2 y of age.